

## MATHEMATICS 360-255-LW

Quantitative Methods II

Martin Huard

Fall 2009

# XX – Inferences about the Population Variance SOLUTIONS

1. The manufacturer of a certain brand of light bulbs claims that the variance of the lives of these bulbs is 4000 square hours. A consumer agency took a random sample of 25 such bulbs and tested them. The variance of the lives of these bulbs was found to be 4990 square hours. Assume that the lives of all such bulbs are (approximately) normally distributed.

- a) Make the 99% confidence interval for the variance and standard deviation of the lives of all such bulbs.

Step 1 Assumptions: The population is normally distributed

Step 2 a) Test statistic:  $\chi^2$  with  $df = 25 - 1 = 24$

b) Level of confidence:  $1 - \alpha = 0.99$  or  $\alpha = 0.01$

Step 3 Point estimate:  $s^2 = 4990$  hours<sup>2</sup>

Step 4 a)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(24, 0.995)} = 9.89$

$$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(24, 0.005)} = 45.56$$


$$\text{b) } \frac{(n-1)s^2}{\chi^2_{(df, \frac{\alpha}{2})}} < \sigma^2 < \frac{(n-1)s^2}{\chi^2_{(df, 1-\frac{\alpha}{2})}}$$

$$\frac{24 \cdot 4990}{45.56} < \sigma^2 < \frac{6 \cdot 4990}{9.89}$$

$$2628 < \sigma^2 < 12109$$

$$51.3 < \sigma < 110.0$$

Step 5 The 99% confidence interval for the variance of the life of light bulbs is 2628 hours<sup>2</sup> to 12109 hours<sup>2</sup>, and for the standard deviation 51.3 hours to 110.0 hours.



- b) Test at the 1% significance level if the variance of such bulbs is different from 4000 square hours. Try with both approaches, the classical and the  $p$ -value.

**Classical Approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 4000 \text{ hours}^2$

$H_A: \sigma^2 \neq 4000 \text{ hours}^2$

Step 3 a) Test statistic:  $\chi^2$  with  $df = 25 - 1 = 24$

b) Two-tailed test with  $\alpha = 0.01$

c)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(24, 0.995)} = 9.89$


$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(24, 0.005)} = 45.56$

Step 4 
$$\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{24 \cdot 4990}{4000} = 29.94$$

Step 5 a)  $\chi^2$  is not in the critical region

b) Fail to reject  $H_0$ .

$\therefore$  There is not sufficient evidence at the 1% level of significance to conclude that the variance of light bulbs is different from 4000 squared hours.



**$p$ -value approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 4000 \text{ hours}^2$

$H_A: \sigma^2 \neq 4000 \text{ hours}^2$

Step 3 a) Test statistic:  $\chi^2$  with  $df = 25 - 1 = 24$

b) Two-tailed test with  $\alpha = 0.01$

Step 4 
$$\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{24 \cdot 4990}{4000} = 29.94$$

$2 \cdot 0.100 < p\text{-value} < 2 \cdot 0.250$

$0.200 < p\text{-value} < 0.500$

Step 5 a)  $p\text{-value} > 0.200 > \alpha = 0.01$

b) Fail to reject  $H_0$ .

$\therefore$  There is not sufficient evidence at the 1% level of significance to conclude that the variance of light bulbs is different from 4000 squared hours.

2. The following are the prices of the same brand of camcorder found at eight stores in Montreal.

\$749 815 789 799 732 825 799 769

- a) Make the 95% confidence interval for the population standard deviation.

Step 1 Assumptions: The population is normally distributed

Step 2 a) Test statistic:  $\chi^2$  with  $df = 8 - 1 = 7$

b) Level of confidence:  $1 - \alpha = 0.95$  or  $\alpha = 0.05$

Step 3 Point estimate:  $s^2 = 1038.3$  squared dollars

Step 4 a)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(7, 0.975)} = 1.69$

$$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(7, 0.025)} = 16.01$$


$$b) \frac{(n-1)s^2}{\chi^2_{(df, \frac{\alpha}{2})}} < \sigma^2 < \frac{(n-1)s^2}{\chi^2_{(df, 1-\frac{\alpha}{2})}}$$

$$\frac{7 \cdot 1038.3}{16.01} < \sigma^2 < \frac{7 \cdot 1038.3}{1.69}$$

$$454.0 < \sigma^2 < 4300.5$$

$$21.31 < \sigma < 65.58$$

Step 5 The 95% confidence interval for the standard deviation of prices of camcorder in Montreal is \$21.31 to \$65.58.



- b) Test at the 5% significance level if the variance of the population is different from 500 square dollars. Use the classical approach. Find the  $p$ -value.

**Classical Approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 500$  squared dollars

$H_A: \sigma^2 \neq 500$  squared dollars

Step 3 a) Test statistic:  $\chi^2$  with  $df = 8 - 1 = 7$

b) Two-tailed test with  $\alpha = 0.05$

c)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(7, 0.975)} = 1.69$


$$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(7, 0.025)} = 16.01$$

Step 4  $\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{7 \cdot 1038.3}{500} = 14.54$

Step 5 a)  $\chi^2$  is not in the critical region

b) Fail to reject  $H_0$ .

$\therefore$  There is not sufficient evidence at the 5% level of significance to conclude that the variance for prices of camcorder in Montreal is different from 500 squared dollars.



**p-value approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 500$  squared dollars

$H_A: \sigma^2 \neq 500$  squared dollars

Step 3 a) Test statistic:  $\chi^2$  with  $df = 8 - 1 = 7$

b) Two-tailed test with  $\alpha = 0.05$

Step 4 
$$\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{7 \cdot 1038.3}{500} = 14.54$$

$2 \cdot 0.025 < p\text{-value} < 2 \cdot 0.050$

$0.050 < p\text{-value} < 0.100$

Step 5 a)  $p\text{-value} > \alpha = 0.05$

b) Fail to reject  $H_0$ .

$\therefore$  There is not sufficient evidence at the 5% level of significance to conclude that the variance for prices of camcorder in Montreal is different from 500 squared dollars.

3. In 1941, the population variance for the age (in years) of a rural Quebec woman at the time of her first marriage was approximately 5.1. A recent study of age of first marriage for a random sample of 41 women in rural Quebec gave a sample variance of 3.3.

- a) Make the 90% confidence interval for the variance and standard deviation for the recent age of first marriage of women in rural Quebec.

Step 1 Assumptions: The population is normally distributed

Step 2 a) Test statistic:  $\chi^2$  with  $df = 41 - 1 = 40$

b) Level of confidence:  $1 - \alpha = 0.90$  or  $\alpha = 0.10$

Step 3 Point estimate:  $s^2 = 3.3$  years<sup>2</sup>

Step 4 a)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(40, 0.95)} = 26.51$

$$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(40, 0.05)} = 55.76$$


b) 
$$\frac{(n-1)s^2}{\chi^2_{(df, \frac{\alpha}{2})}} < \sigma^2 < \frac{(n-1)s^2}{\chi^2_{(df, 1-\frac{\alpha}{2})}}$$

$$\frac{40 \cdot 3.3}{55.76} < \sigma^2 < \frac{40 \cdot 3.3}{26.51}$$

$$2.367 < \sigma^2 < 4.979$$

$$1.54 < \sigma < 2.23$$

Step 5 The 90% confidence interval for the variance for the age of first marriage of woman in rural Quebec is 2.367 years<sup>2</sup> to 4.979 years<sup>2</sup>, and for the standard deviation is 1.54 years to 2.23 years.



- b) Test at the 10% significance level the claim that the current variance is less than 5.1. Use the classical approach. Find the  $p$ -value.

**Classical Approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 5.1 \text{ years}^2$

$H_A: \sigma^2 < 5.1 \text{ years}^2$

Step 3 a) Test statistic:  $\chi^2$  with  $df = 41 - 1 = 40$

b) Left-tailed test with  $\alpha = 0.10$


c)  $\chi^2_{(df, 1-\alpha)} = \chi^2_{(40, 0.90)} = 29.05$

Step 4  $\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{40 \cdot 3.3}{5.1} = 25.88$

Step 5 a)  $\chi^2$  is in the critical region

b) Reject  $H_0$ .

$\therefore$  There is sufficient evidence at the 10% level of significance to conclude that the current variance for the age of first marriage of woman in rural Quebec is less than 5.1



**$p$ -value approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 5.1 \text{ years}^2$

$H_A: \sigma^2 < 5.1 \text{ years}^2$

Step 3 a) Test statistic:  $\chi^2$  with  $df = 41 - 1 = 40$

b) Left-tailed test with  $\alpha = 0.10$

Step 4  $\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{40 \cdot 3.3}{5.1} = 25.88$

$1 - 0.975 < p\text{-value} < 1 - 0.950$

$0.025 < p\text{-value} < 0.05$

Step 5 a)  $p\text{-value} < 0.05 < \alpha = 0.10$

b) Reject  $H_0$ .

$\therefore$  There is sufficient evidence at the 10% level of significance to conclude that the current variance for the age of first marriage of woman in rural Quebec is less than 5.1

4. A machine that puts corn flakes in boxes is adjusted to put an average of 400 grams in each box with a standard deviation of 6 grams. A random sample of 12 boxes gave a sample standard deviation of 9 grams.

- a) Make the 95% confidence interval for the standard deviation for the volume in corn flakes boxes.

Step 1 Assumptions: The population is normally distributed

Step 2 a) Test statistic:  $\chi^2$  with  $df = 12 - 1 = 11$

b) Level of confidence:  $1 - \alpha = 0.95$  or  $\alpha = 0.05$

Step 3 Point estimate:  $s^2 = 9^2 = 81$  grams<sup>2</sup>

Step 4 a)  $\chi^2_{(df, 1-\frac{\alpha}{2})} = \chi^2_{(11, 0.975)} = 3.82$

$$\chi^2_{(df, \frac{\alpha}{2})} = \chi^2_{(11, 0.025)} = 21.92$$


$$b) \frac{(n-1)s^2}{\chi^2_{(df, \frac{\alpha}{2})}} < \sigma^2 < \frac{(n-1)s^2}{\chi^2_{(df, 1-\frac{\alpha}{2})}}$$

$$\frac{11 \cdot 81}{21.92} < \sigma^2 < \frac{11 \cdot 81}{3.82}$$

$$40.65 < \sigma^2 < 233.25$$

$$6.38 < \sigma < 15.27$$

Step 5 The 95% confidence interval for the standard deviation for the volume in corn flakes boxes is 6.38 grams to 15.27 grams.



- b) Do these data support the claim that the variance has increased and the machine needs to be brought back into adjustment? Use the classical approach. Find the  $p$ -value.

**Classical Approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 6^2 = 36$  grams<sup>2</sup>

$H_A: \sigma^2 > 36$  grams<sup>2</sup>

Step 3 a) Test statistic:  $\chi^2$  with  $df = 12 - 1 = 11$

b) Left-tailed test with  $\alpha = 0.05$


c)  $\chi^2_{(df, \alpha)} = \chi^2_{(11, 0.05)} = 19.68$

Step 4  $\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{11 \cdot 81}{36} = 24.75$

Step 5 a)  $\chi^2$  is in the critical region

b) Reject  $H_0$ .

$\therefore$  There is sufficient evidence at the 5 level of significance to conclude that the machine needs to be adjusted



***p*-value approach**

Step 1 Assumptions: The population is normally distributed

Step 2  $H_0: \sigma^2 = 6^2 = 36 \text{ grams}^2$

$H_A: \sigma^2 > 36 \text{ grams}^2$

Step 3 a) Test statistic:  $\chi^2$  with  $df = 12 - 1 = 11$

b) Left-tailed test with  $\alpha = 0.05$

Step 4 
$$\chi^2 = \frac{(n-1)s^2}{\sigma^2} = \frac{11 \cdot 81}{36} = 24.75$$

$0.005 < p\text{-value} < 0.01$

Step 5 a)  $p\text{-value} < 0.01 < \alpha = 0.05$

b) Reject  $H_0$ .

$\therefore$  There is sufficient evidence at the 5 level of significance to conclude that the machine needs to be adjusted.